

[Cat. No.] Please refer to the **Ordering Information**

Introduction

AccuPower® RocketScript™ RT-PCR PreMix, RNase H Minus is a one-step RT-PCR product that uses RocketScript™ Reverse Transcriptase (RTase), RNase H Minus and ProFi Taq DNA Polymerase independently developed by BIONEER. RocketScript™ RTase, RNase H Minus is a recombinant *M-MLV* RTase with enhanced thermostability up to 70°C, as well as eliminated RNase H activity. It can produce efficiently full-length cDNA from long RNA transcripts up to 12.5 kb and synthesize cDNA even with small amounts of 1 pg of total RNA. In addition, included ProFi Taq DNA Polymerase provides accurate amplification and higher fidelity for PCR. This product contains vacuum-dried components including RocketScript™ RTase, reaction buffer, DTT, dNTPs, RNase inhibitor, and ProFi Taq DNA Polymerase. It simplifies preparation of reverse transcription reaction mixture and PCR mixture by adding template RNA and primers without any extra process.




Figure 1. Elimination of RNase H activity. Eliminated RNase H activity from RTase complex enables the RTase to produce longer RNA transcripts, up to 12.5 kb, compared with original RocketScript™ RTase itself.

Applications

- Standard RT-PCR
- Long kb RT-PCR
- Virus detection
- Gene expression analysis

Features & Benefits

- **Thermostable activity:** RocketScript™ RTase has excellent thermal resistance even at 70°C, so it is suitable for cDNA synthesis of secondary RNA structures.
- **No RNase H activity:** Useful for cDNA synthesis up to 12.5 kb from long target RNA using enzymes developed by proprietary genetic engineering technology.
- **User-friendly:** Simple amplification of long-size RNA difficult to be done with conventional one-step RT-PCR methods.

Composition

Composition	Concentration
RocketScript™ Reverse Transcriptase, RNase H Minus	200 U
5X Reaction buffer	1X
DTT	0.25 mM
dNTPs (dATP, dCTP, dGTP, dTTP)	Each 250 µM
RNase inhibitor	1 U
ProFi Taq DNA Polymerase	1 U
Stabilizer and tracking dye	1X

Specifications

RocketScript™ Reverse Transcriptase, RNase H Minus	
DNase activity	No
RNase activity	No
RNase H activity	No
Fragment size	Up to 12.5 kb

Storage

Store at -20°C. If stored in the recommended temperature, this product will be stable until the expiration date printed out on the label.

Online Resources



Korean



English

Visit our **product page** for additional information and protocols


Ordering Information

Description	Cat. No.
0.2 ml thin-wall 96 tubes	K-2231
8-tube strips	K-2233
with attached cap	K-2232
480 tubes	K-2234


Notice

BIONEER corporation reserves the right to make corrections, modifications, improvements and other changes to its products, services, specifications or product descriptions at any time without notice.

Explanation of Symbols



Experimental Procedures

Steps		Procedure Details																												
1	 Preparation of reaction mixture	<p>1. Add template RNA, primers and nuclease-free water into <i>AccuPower® RocketScript™</i> RT-PCR PreMix, RNase H Minus tubes to make a total volume of 20 µl or 50 µl. Do not include the dried pellet.</p> <ul style="list-style-type: none"> Amount of template RNA and primers <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th colspan="2">Components</th> <th>20 µl reaction</th> <th>50 µl reaction</th> </tr> </thead> <tbody> <tr> <td rowspan="2">Template RNA</td> <td>Total RNA</td> <td>0.01-5 µg</td> <td>0.01-5 µg</td> </tr> <tr> <td>Poly(A) RNA</td> <td>0.01-5 µg</td> <td>0.01-5 µg</td> </tr> <tr> <td>Primers</td> <td>Gene specific primer</td> <td>10-30 pmol</td> <td>10-50 pmol</td> </tr> </tbody> </table> <p>2. Dissolve the vacuum-dried pellet by pipetting or vortexing, and briefly spin down.</p>	Components		20 µl reaction	50 µl reaction	Template RNA	Total RNA	0.01-5 µg	0.01-5 µg	Poly(A) RNA	0.01-5 µg	0.01-5 µg	Primers	Gene specific primer	10-30 pmol	10-50 pmol													
Components		20 µl reaction	50 µl reaction																											
Template RNA	Total RNA	0.01-5 µg	0.01-5 µg																											
	Poly(A) RNA	0.01-5 µg	0.01-5 µg																											
Primers	Gene specific primer	10-30 pmol	10-50 pmol																											
2	 RT-PCR	<p>3. Perform the reaction under the following conditions.</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Step</th> <th>Temperature</th> <th>Time</th> <th>Cycles</th> </tr> </thead> <tbody> <tr> <td>cDNA synthesis</td> <td>42-70°C</td> <td>30 min</td> <td>1 cycle</td> </tr> <tr> <td>Pre-denaturation</td> <td>95°C</td> <td>5 min</td> <td>1 cycle</td> </tr> <tr> <td>Denaturation</td> <td>95°C</td> <td>10-30 sec</td> <td></td> </tr> <tr> <td>Annealing</td> <td>55°C</td> <td>10-30 sec</td> <td>30-35 cycles</td> </tr> <tr> <td>Extension</td> <td>72°C</td> <td>1 kb/min</td> <td></td> </tr> <tr> <td>Final extension</td> <td>72°C</td> <td>5 min</td> <td>1 cycle</td> </tr> </tbody> </table> <p>* Note: Reaction temperature should be optimized according to T_m value of primers.</p>	Step	Temperature	Time	Cycles	cDNA synthesis	42-70°C	30 min	1 cycle	Pre-denaturation	95°C	5 min	1 cycle	Denaturation	95°C	10-30 sec		Annealing	55°C	10-30 sec	30-35 cycles	Extension	72°C	1 kb/min		Final extension	72°C	5 min	1 cycle
Step	Temperature	Time	Cycles																											
cDNA synthesis	42-70°C	30 min	1 cycle																											
Pre-denaturation	95°C	5 min	1 cycle																											
Denaturation	95°C	10-30 sec																												
Annealing	55°C	10-30 sec	30-35 cycles																											
Extension	72°C	1 kb/min																												
Final extension	72°C	5 min	1 cycle																											
3	 Analyze with gel electrophoresis	<p>4. After the reaction, maintain the reaction mixture at 4°C. The samples can be stored at -20°C until use.</p> <p>5. Load samples on agarose gel without adding a loading-dye mixture, and perform gel electrophoresis for analysis.</p>																												