

Bioneer Corporation
8-11 Munpyeongseo-ro, Daedeok-gu, Daejeon, 34302
Republic of Korea
Tel: +82-42-930-8777 (Korea : 1588-9788)
Fax: +82-42-930-8688 E-mail: sales@bioneer.com

Bioneer Inc.
155 Filbert St. Suite 216
Oakland, CA 94607, USA
Toll Free: +1-877-264-4300 Fax: +1-510-865-0350
E-mail: order.usa@bioneer.us.com

Bioneer R&D Center
Korea Bio Park BLDG #B-702, 700 Daewangpangyo-ro
Bundang-gu, Seongnam-si, Gyeonggi-do, 13488
Republic of Korea
Tel: +82-31-628-0500 Fax: +82-31-628-0555

AccuPower® RT-PCR PreMix contains all the components necessary for cDNA synthesis and amplification in one tube.

Advantages

- Speed** Substantial reduction in reaction setup time.
- Stability** As each tube of *AccuPower®* RT-PCR PreMix contains a stabilizer (patented in US and Korea), which can maintain the stability of the RTase and DNA polymerase up to 2 years at -20°C. Pre-denaturation during PCR has no effects on the activity of the DNA polymerase.
- Reproducibility and Yield** The strict functional QC assays demonstrated highly consistent and reproducible RT/PCR performance. In most applications an increase in yield is observed as compared to the standard reactions.
- Simplicity** The fewer manual steps allow reduction in potential errors. Each tube contains tracking dye and precipitant for agarose gel electrophoresis, eliminating the needs for a separate loading buffer.

Experimental Protocol

- Mix the template RNA and the reverse primer in a sterile tube as indicated below;

Concentration of template RNA and primer			
Reaction volume		20 µl reaction	50 µl reaction
Template RNA	Total RNA	0.5~1.0 µg	1.0~2.0 µg
	Poly(A) RNA	0.05~0.1 µg	0.1~0.2 µg
Primer	Primer	10~30 pmole	20~50 pmole

- Incubate the mixture at 70°C for 5 min and place it on ice.
- Transfer the incubated mixture and the forward primer to *AccuPower®* RT/PCR PreMix tube, and then fill up the reaction volume with DEPC-DW.
- Dissolve the vacuum dried blue pellet by vortexing, and briefly spin down.
- Add mineral oil to each tube (This step is unnecessary when using a thermal cycler with top heating.).
- Perform cDNA synthesis reaction as follows :
 - 42°C, 60 min. (cDNA synthesis)
 - 94°C, 5 min. (RTase inactivation)
- Perform PCR cycles according to the PCR condition.
(Annealing temperature and time need to be optimized for each primer/template combination.)

Ordering Information

Tube type	Reaction	Cat.No	Description	Tube type	Reaction	Cat.No	Description
0.2 ml Tube	20 µl	K-2055	0.2 ml thin-wall 8-strip tubes with attached cap / 96 tubes	96-well	20 µl	K-2262-4	thin-wall 96-well flat plate
		K-2055-B	0.2 ml thin-wall 8-strip tubes with attached cap / 480 tubes			K-2262-5	thin-wall 96-well full-skirted plate
	50 µl	K-2057	0.2 ml thin-wall 8-strip tubes with attached cap / 96 tubes			K-2262-6	thin-wall 96-well semi-skirted plate
		K-2057-B	0.2 ml thin-wall 8-strip tubes with attached cap / 480 tubes	K-2084-1	thin-wall 384-well full-skirted plate		
96-well	10 µl	K-2262-1	thin-wall 96-well flat plate	384-well	10 µl	K-2084-2	thin-wall 384-well full-skirted plate
		K-2262-2	thin-wall 96-well full-skirted plate		20 µl	K-2084-3	thin-wall 384-well full-skirted plate
		K-2262-3	thin-wall 96-well semi-skirted plate		0.5 ml Tube	50 µl	k-2056

Notice

Bioneer corporation reserves the right to make corrections, modifications, improvements and other changes to its products, services, specifications or product descriptions at any time without notice. All information provided here is subject to change without notice.

A complete product list appears on our web site at www.bioneer.com.